

Issues in Specific Clinical Situations

1. Amin RM, Puvanesarajah V, Chaudhry YP, et al. Reducing unnecessary crossmatching for hip fracture patients by accounting for preoperative hemoglobin concentration. *World J Orthop* 2021;12(5):292-300. <https://doi.org/10.5312/wjo.v12.i5.292>. PMID: 34055586; PMCID: PMC8152439.
2. Winearls J, Reade MC, McQuilten Z, Curry N. Fibrinogen in traumatic haemorrhage. *Curr Opin Anaesthesiol* 2021;34(4):514-20. <https://doi.org/10.1097/aco.0000000000001027>. PMID: 34091461.
3. Amanvermez Senarslan D, Yildirim F, Kurdal AT, et al. Efficacy and cost-effectiveness of cell saver usage in the repair of thoracic aortic aneurysms and dissections. *Perfusion* 2021; 2676591211028178. <https://doi.org/10.1177%2F02676591211028178>. PMID: 34192996.
4. Walczak S, Velanovich V. Prediction of perioperative transfusions using an artificial neural network. *PLoS One* 2020;15(2):e0229450. <https://doi.org/10.1371/journal.pone.0229450>. PMID: 32092108; PMCID: PMC7039514.
5. Marcondes SS, Carrareto AR, Zago-Gomes MDP, et al. Evaluation of the use of blood in surgeries as a tool to change patterns for requesting blood product reserves. *Clinics (Sao Paulo)*. 2019;74:e652. <https://doi.org/10.6061/clinics/2019/e652>. PMID: 31038643; PMCID: PMC6467177.
6. Hallet J, Mahar AL, Nathens AB, et al. The impact of perioperative blood transfusions on short-term outcomes following hepatectomy. *Hepatobiliary Surg Nutr* 2018;7(1):1-10. <https://doi.org/10.21037/hbsn.2017.05.07>. PMID: 29531938; PMCID: PMC5835602.
7. Bisbe E, Basora M, Colomina MJ. Spanish Best Practice in Peri-operative Anaemia Optimisation Panel. Peri-operative treatment of anaemia in major orthopaedic surgery: A practical approach from Spain. *Blood Transfus* 2017;15(4):296-306. <https://doi.org/10.2450/2017.0177-16>. PMID: 28151388; PMCID: PMC5490724.
8. O'Donnell C, Michael N, Bloch N, et al. Strategies to minimize blood loss and transfusion in pediatric spine surgery. *JBJS Rev* 2017;5(5):e1. <https://doi.org/10.2106/jbjs.rvw.16.00064>. PMID: 28471775.
9. Arias-Morales CE, Stoicea N, Gonzalez-Zacarias AA, et al. Revisiting blood transfusion and predictors of outcome in cardiac surgery patients: A concise perspective. *F1000Res* 2017;6 (F1000 Faculty Rev):168. <https://doi.org/10.12688/f1000research.10085.1>. PMID: 28299184; PMCID: PMC5321117.
10. Woodrum CL, Wisniewski M, Triulzi DJ, et al. The effects of a data driven maximum surgical blood ordering schedule on preoperative blood ordering practices. *Hematology* 2017;22(9):571-7. <https://doi.org/10.1080/10245332.2017.1318336>. PMID: 28441911.
11. Elmi M, Mahar A, Kagedan D, et al. The impact of blood transfusion on perioperative outcomes following gastric cancer resection: An analysis of the American College of Surgeons National Surgical Quality Improvement Program database. *Can J Surg* 2016;59(5):322-9. <https://doi.org/10.1503/cjs.004016>. PMID: 27668330; PMCID: PMC5042719.
12. Yohanathan L, Coburn NG, McLeod RS, et al. Best Practices in general surgery (BPIGS) group. Understanding perioperative transfusion practices in gastrointestinal surgery—a prac-

- tice survey of general surgeons. *J Gastrointest Surg* 2016;20(6):1106-22. <https://doi.org/10.1007/s11605-016-3111-15>. PMID: 27025709.
13. Goobie SM, Haas T. Perioperative bleeding management in pediatric patients. *Curr Opin Anaesthesiol* 2016;29(3):352-8. <https://doi.org/10.1097/ACO.0000000000000308>. PMID: 26844864.
 14. Rinehart JB, Lee TC, Kaneshiro K, et al. Perioperative blood ordering optimization process using information from an anesthesia information management system. *Transfusion* 2016;56(4):938-45. <https://doi.org/10.1111/trf.13492>. PMID: 26876784.
 15. Hallet J, Mahar AL, Tsang ME, et al. The impact of peri-operative blood transfusions on post-pancreatectomy short-term outcomes: An analysis from the American College of Surgeons National Surgical Quality Improvement Program. *HPB (Oxford)* 2015;17(11):975-82. <https://doi.org/10.1111/hpb.12473>. PMID: 26301741; PMCID: PMC4605335.
 16. White MJ, Hazard SW 3rd, Frank SM, et al. The evolution of perioperative transfusion testing and blood ordering. *Anesth Analg* 2015;120(6):1196-203. <https://doi.org/10.1213/ANE.0000000000000619>. PMID: 25988630.
 17. Pasternak J, Nikolic D, Milosevic D, et al. An analysis of the influence of intra-operative blood salvage and autologous transfusion on reducing the need for allogeneic transfusion in elective infrarenal abdominal aortic aneurysm repair. *Blood Transfus* 2014;12(Suppl 1):s182-s186. <https://doi.org/10.2450/2012.0069-12>. PMID: 23114525; PMCID: PMC3934265.

Volume Resuscitation

1. van den Brink DP, Kleinveld DJB, Sloos PH, et al. Plasma as a resuscitation fluid for volume-depleted shock: Potential benefits and risks. *Transfusion* 2021;61(Suppl 1):S301-S312. <https://doi.org/10.1111/trf.16462>. PMID: 34057210.
2. Walsh M, Moore EE, Moore HB, et al. Whole blood, fixed ratio, or goal-directed blood component therapy for the initial resuscitation of severely hemorrhaging trauma patients: A narrative review. *J Clin Med* 2021;10(2):320. <https://doi.org/10.3390/jcm10020320>. PMID: 33477257; PMCID: PMC7830337.
3. Munoz JL, Kimura AM, Xenakis E, et al. Whole blood transfusion reduces overall component transfusion in cases of placenta accreta spectrum: A pilot program. *J Matern Fetal Neonatal Med* 2021:1-6. <https://doi.org/10.1080/14767058.2021.1915275>. PMID: 33902384.
4. Nowadly CD, Fisher AD, Borgman MA, et al. The use of whole blood transfusion during non-traumatic resuscitation. *Mil Med* 2021:usab128. <https://doi.org/10.1093/milmed/usab128>. PMID: 33856481.
5. Salamea-Molina JC, Himmler AN, Valencia-Angel LI, et al. Whole blood for blood loss: Hemostatic resuscitation in damage control. *Colomb Med (Cali)* 2020;51(4):e4044511. <https://doi.org/10.25100/cm.v51i4.4511>. PMID: 33795899; PMCID: PMC7968429.
6. Malkin M, Nevo A, Brundage SI, Schreiber M. Effectiveness and safety of whole blood compared to balanced blood components in resuscitation of hemorrhaging trauma patients - A systematic review. *Injury* 2021;52(2):182-8. <https://doi.org/10.1016/j.injury.2020.10.095>. PMID: 33160609.
7. Leeper CM, Yazer MH, Triulzi DJ, et al. Whole blood is superior to component transfusion for injured children: A propensity matched analysis. *Ann Surg* 2020;272(4):590-4. <https://doi.org/10.1097/SLA.0000000000004378>. PMID: 32932312.

8. Polites SF, Moody S, Williams RF, et al. Timing and volume of crystalloid and blood products in pediatric trauma: An eastern association for the surgery of trauma multicenter prospective observational study. *J Trauma Acute Care Surg* 2020;89(1):36-42. <https://doi.org/10.1097/TA.0000000000002702>. PMID: 32251263.
9. Anto VP, Guyette FX, Brown J, et al and the PAMPer study group. Severity of hemorrhage and the survival benefit associated with plasma: Results from a randomized prehospital plasma trial. *J Trauma Acute Care Surg* 2020;88(1):141-7. <https://doi.org/10.1097/TA.0000000000002530>. PMID: 31688793.
10. Hazelton JP, Cannon JW, Zatorski C, et al. Cold-stored whole blood: A better method of trauma resuscitation? *J Trauma Acute Care Surg* 2019;87(5):1035-41. <https://doi.org/10.1097/TA.0000000000002471>. PMID: 31389912.
11. Geeraedts LM Jr, Pothof LA, Caldwell E, et al. Prehospital fluid resuscitation in hypotensive trauma patients: Do we need a tailored approach? *Injury* 2015;46(1):4-9. <https://doi.org/10.1016/j.injury.2014.08.001>. PMID: 25173671.
12. Magnotti LJ, Zarzaur BL, Fischer PE, et al. Improved survival after hemostatic resuscitation: Does the emperor have no clothes? *J Trauma* 2011;70(1):97-102. <https://doi.org/10.1097/TA.0b013e3182051691>. PMID: 21217487.

Cardiology

1. Carson JL, Sieber F, Cook DR, et al. Liberal versus restrictive blood transfusion strategy: 3-year survival and cause of death results from the FOCUS randomised controlled trial. *The Lancet* 2015;385(9974):1183-9. [https://doi.org/10.1016/s0140-6736\(14\)62286-8](https://doi.org/10.1016/s0140-6736(14)62286-8).
2. Bolliger D, Mauermann E, Tanaka KA. Thresholds for perioperative administration of hemostatic blood components and coagulation factor concentrates: An unmet medical need. *J Cardiothorac Vasc Anesth* 2015;29(3):768-76. <https://doi.org/10.1053/j.jvca.2015.02.023>.
3. Geissler RG, Rotering H, Buddendick H, et al. Utilisation of blood components in cardiac surgery: A single-centre retrospective analysis with regard to diagnosis-related procedures. *Transfus Med Hemother* 2015;42(2):75-82. <https://doi.org/10.1159/000377691>.
4. Mangwana S, Bedi N, Yadav P, Chugh R. Optimization of blood transfusion services: Analysis of blood requisition and utilization practices in cardiac surgical patients in a tertiary care hospital, India. *Glob J Transfus Med* 2017;2:47-51. <https://doi.org/10.4103/2468-8398.202713>.
5. Meybohm P, Richards T, Isbister J, et al. Patient blood management bundles to facilitate implementation. *Transfus Med Rev* 2016;31(1):62-71. <https://doi.org/10.1016/j.tmr.2016.05.012>.
6. Hensley NB, Kostibas MP, Yang W, et al. Blood utilization in revision versus first-time cardiac surgery: An update in the era of patient blood management. *Transfusion* 2017;58(1):168-75. <https://doi.org/10.1111/trf.14361>.
7. Boer C, Meesters MI, Milojevic M, et al. 2017 EACTS/EACTA. Guidelines on patient blood management for adult cardiac surgery. *J Cardiothorac Vasc Anesth* 2018;32(1):88-120. <https://doi.org/10.1053/j.jvca.2017.06.026>.